

UCF



Stands For Opportunity

---

---

*CDA6530: Performance Models of Computers and Networks*

***Chapter 9: Statistical Analysis of  
Simulated Data and Confidence Interval***

# Sample Mean

- r.v.  $X$ :  $E[X]=\theta$ ,  $\text{Var}[X]=\sigma^2$
- Q: how to use simulation to derive?
  - Simulate  $X$  repeatedly
- $X_1, \dots, X_n$  are i.i.d., =<sub>statistic</sub>  $X$
- Sample mean:  $\bar{X} \equiv \sum_{i=1}^n \frac{X_i}{n}$

$$E[\bar{X}] = \theta \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n}$$

# Sample Variance

- $\sigma^2$  unknown in simulation
  - Hard to use  $Var(\bar{X}) = \frac{\sigma^2}{n}$  to measure simulation variance
  - Thus we need to estimate  $\sigma^2$

- **Sample variance  $S^2$ :**

$$S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n - 1}$$

- n-1 instead of n is to provide unbiased estimator  $E[S^2] = \sigma^2$

# Estimate Error

---

- Sample mean  $\bar{X}$  is a good estimator of  $\theta$ , but has an error
  - How confident we are sure that the sample mean is within an acceptable error?
- From central limit theorem:

$$\sqrt{n} \frac{(\bar{X} - \theta)}{\sigma} \sim N(0, 1)$$

- It means that:

$$\sqrt{n} \frac{(\bar{X} - \theta)}{S} \sim N(0, 1)$$

# Confidence Interval

- R.v.  $Z \sim N(0,1)$ , for  $0 < \alpha < 1$ , define:
  - $P(Z > z_\alpha) = \alpha$
  - $z_{0.025} = 1.96$
  - $P(-z_{0.025} < Z < z_{0.025}) = 1 - 2\alpha = 0.95$

$$P\left(\bar{X} - z_{\alpha/2} \frac{S}{\sqrt{n}} < \theta < \bar{X} + z_{\alpha/2} \frac{S}{\sqrt{n}}\right) \approx 1 - \alpha$$

$$P\left(\bar{X} - 1.96 \frac{S}{\sqrt{n}} < \theta < \bar{X} + 1.96 \frac{S}{\sqrt{n}}\right) \approx 0.95$$

95% confidence interval ( $\alpha = 0.05$ ) of an estimate is:

$$\left(\bar{X} \pm 1.96 S / \sqrt{n}\right)$$

# When to stop a simulation?

---

- Repeatedly generate data (sample) until 100(1- $\alpha$ ) percent confidence interval estimate of  $\theta$  is less than  $I$ 
  - Generate at least 100 data values.
  - Continue generate, until you generated  $k$  values such that  $2z_{\alpha/2}S/\sqrt{k} < I$
  - The 100(1- $\alpha$ ) percent confidence interval of estimate is


$$(\bar{X} - z_{\alpha/2}S/\sqrt{k}, \bar{X} + z_{\alpha/2}S/\sqrt{k})$$

# Fix no. of simulation runs


- Suppose we only simulate 100 times
  - $k=100$
- What is the 95% confidence interval?

$$(\bar{X} - z_{\alpha/2}S/\sqrt{k}, \bar{X} + z_{\alpha/2}S/\sqrt{k})$$

$$(\bar{X} - 1.96S/\sqrt{k}, \bar{X} + 1.96S/\sqrt{k})$$



# Example: Generating Expo. Distribution



Compare 100 samples and 1000 samples confidence intervals